



KCNQ3 gene

potassium voltage-gated channel subfamily Q member 3

Normal Function

The *KCNQ3* gene belongs to a large family of genes that provide instructions for making potassium channels. These channels, which transport positively charged atoms (ions) of potassium into and out of cells, play a key role in a cell's ability to generate and transmit electrical signals.

The specific function of a potassium channel depends on its protein components and its location in the body. Channels made with the KCNQ3 protein are active in nerve cells (neurons) in the brain, where they transport potassium ions out of cells. These channels transmit a particular type of electrical signal called the M-current, which prevents the neuron from continuing to send signals to other neurons. The M-current ensures that the neuron is not constantly active, or excitable.

Potassium channels are made up of several protein components (subunits). Each channel contains four alpha subunits that form the hole (pore) through which potassium ions move. Four alpha subunits from the *KCNQ3* gene can form a channel. However, the KCNQ3 alpha subunits can also interact with alpha subunits from the *KCNQ2* gene to form a functional potassium channel, and these channels transmit a much stronger M-current.

Health Conditions Related to Genetic Changes


benign familial neonatal seizures

A mutation in the *KCNQ3* gene has been identified in some people with benign familial neonatal seizures (BFNS), a condition characterized by recurrent seizures in newborn babies. The seizures begin around day 3 of life and usually go away within 1 to 4 months. At least three mutations have been identified in people with this condition, and these mutations change single protein building blocks (amino acids) in the KCNQ3 protein. As a result of these mutations, the M-current is reduced. Researchers believe that a reduction of the current by 25 percent is enough to cause BFNS. A reduced M-current leads to excessive excitability of neurons, which is known to cause seizures. It is unclear why the seizures stop around the age of 4 months. It has been suggested that potassium channels formed from the KCNQ2 and KCNQ3 proteins play a major role in preventing excessive excitability of neurons in newborns, but other mechanisms develop during infancy.

Chromosomal Location

Cytogenetic Location: 8q24.22, which is the long (q) arm of chromosome 8 at position 24.22

Molecular Location: base pairs 132,120,858 to 132,480,757 on chromosome 8 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- BFNC2
- EBN2
- KCNQ3_HUMAN
- KV7.3
- potassium channel subunit alpha KvLQT3
- potassium channel, voltage gated KQT-like subfamily Q, member 3
- potassium channel, voltage-gated, subfamily Q, member 3
- potassium voltage-gated channel subfamily KQT member 3
- potassium voltage-gated channel, KQT-like subfamily, member 3
- voltage-gated potassium channel subunit Kv7.3

Additional Information & Resources

Educational Resources

- Biochemistry (5th Edition, 2002): Action Potentials are Mediated by Transient Changes in Na⁺ and K⁺ Permeability
<https://www.ncbi.nlm.nih.gov/books/NBK22509/#A1816>

GeneReviews

- KCNQ2-Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK32534>
- KCNQ3-Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK201978>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28KCNQ3%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- POTASSIUM CHANNEL, VOLTAGE-GATED, KQT-LIKE SUBFAMILY, MEMBER 3
<http://omim.org/entry/602232>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_KCNQ3.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=KCNQ3%5Bgene%5D>
- HGNC Gene Family: Potassium voltage-gated channels
<http://www.genenames.org/cgi-bin/genefamilies/set/274>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=6297
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/3786>
- UniProt
<http://www.uniprot.org/uniprot/O43525>

Sources for This Summary

- Chung HJ, Jan YN, Jan LY. Polarized axonal surface expression of neuronal KCNQ channels is mediated by multiple signals in the KCNQ2 and KCNQ3 C-terminal domains. *Proc Natl Acad Sci U S A*. 2006 Jun 6;103(23):8870-5. Epub 2006 May 30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16735477>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1472242/>
- OMIM: POTASSIUM CHANNEL, VOLTAGE-GATED, KQT-LIKE SUBFAMILY, MEMBER 3
<http://omim.org/entry/602232>

- Rogawski MA. KCNQ2/KCNQ3 K⁺ channels and the molecular pathogenesis of epilepsy: implications for therapy. *Trends Neurosci.* 2000 Sep;23(9):393-8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10941184>
- Schroeder BC, Kubisch C, Stein V, Jentsch TJ. Moderate loss of function of cyclic-AMP-modulated KCNQ2/KCNQ3 K⁺ channels causes epilepsy. *Nature.* 1998 Dec 17;396(6712):687-90.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9872318>
- Singh NA, Westenskow P, Charlier C, Pappas C, Leslie J, Dillon J, Anderson VE, Sanguinetti MC, Leppert MF; BFNC Physician Consortium. KCNQ2 and KCNQ3 potassium channel genes in benign familial neonatal convulsions: expansion of the functional and mutation spectrum. *Brain.* 2003 Dec; 126(Pt 12):2726-37. Epub 2003 Oct 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14534157>
- Soldovieri MV, Miceli F, Bellini G, Coppola G, Pascotto A, Taglialatela M. Correlating the clinical and genetic features of benign familial neonatal seizures (BFNS) with the functional consequences of underlying mutations. *Channels (Austin).* 2007 Jul-Aug;1(4):228-33. Epub 2007 Aug 2.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18698150>
- Wang HS, Pan Z, Shi W, Brown BS, Wymore RS, Cohen IS, Dixon JE, McKinnon D. KCNQ2 and KCNQ3 potassium channel subunits: molecular correlates of the M-channel. *Science.* 1998 Dec 4; 282(5395):1890-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9836639>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/KCNQ3>

Reviewed: May 2011
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services